



# Lecture 24

---

## Interpreting Confidence

Slides created by John DeNero ([denero@berkeley.edu](mailto:denero@berkeley.edu)) and Ani Adhikari ([adhikari@berkeley.edu](mailto:adhikari@berkeley.edu))

# Announcements


# The Bootstrap

# Key to Resampling

---

- From the original sample,
    - draw at random
    - with replacement
    - as many values as the original sample contained
  - The size of the new sample has to be the same as the original one, so that the two estimates are comparable
-

# Why the Bootstrap Works




# Inference Using the Bootstrap




population



sample



resamples



All of these look pretty similar, most likely.

# 95% Confidence Interval

---

- Interval of **estimates of a parameter**
- Based on random sampling
- 95% is called the confidence level
  - Could be any percent between 0 and 100
  - Higher level means wider intervals
- The **confidence is in the process** that generated the interval:
  - It generates a “good” interval about 95% of the time.

(Demo)

---

**Use Methods Appropriately**



# Can You Use a CI Like This?

---

By our calculation, an approximate 95% confidence interval for the average age of the mothers in the population is (26.9, 27.6) years.

## True or False:

- About 95% of the mothers in the population were between 26.9 years and 27.6 years old.

**Answer: False.** We're estimating that their **average age** is in this interval.

(Demo)

---

# Is This What a CI Means?

---

An approximate 95% confidence interval for the average age of the mothers in the population is (26.9, 27.6) years.

## True or False:

- There is a 0.95 probability that the average age of mothers in the population is in the range 26.9 to 27.6 years.

**Answer: False.** The average age of the mothers in the population is unknown but it's a constant. It's not random. No chances involved.

---

# When *Not* to Use The Bootstrap

---

- If you're trying to estimate very high or very low percentiles, or min and max
- If you're trying to estimate any parameter that's greatly affected by rare elements of the population
- If the probability distribution of your statistic is not roughly bell shaped (the shape of the empirical distribution will be a clue)
- If the original sample is very small

(Demo)

---

# Confidence Intervals For Testing

# Using a CI for Testing

---

- Null hypothesis: **Population average =  $x$**
- Alternative hypothesis: **Population average  $\neq x$**
- Cutoff for P-value:  $p\%$
- Method:
  - Construct a  $(100-p)\%$  confidence interval for the population average
  - If  $x$  is not in the interval, reject the null
  - If  $x$  is in the interval, can't reject the null

(Demo)

---